

Source: [Legal > / ... / > U.S. Patents, European Patents, Patent Abstracts of Japan, PCT Patents, and U.K. !\[\]\(c8d96c8885d3000a912c2582004aed63\_img.jpg\)](#)  
 Terms: [transglutaminase /p \(cross-link! or crosslink!\) and date<8/1/1998](#) ([Edit Search](#) | [Suggest Terms for My Search](#))

 Select for FOCUS™ or Delivery



95934272 745670 June 23, 2004

## EUROPEAN PATENT

745670

[Access PDF of Official Patent \\*](#)

[Check for Patent Family Report PDF availability \\*](#)

\* Note: A transactional charge will be incurred for downloading an Official Patent or Patent Family Report. Your acceptance of this charge occurs in a later step in your session. The transactional charge for downloading is outside of customer subscriptions; it is not included in any flat rate packages.

[Order Patent File History / Wrapper from REEDFAX®](#)  
[Link to Claims Section](#)

**December 4, 1996, Issue No.: 199649**

STABILIZED TRANSGLUTAMINASE AND ENZYMATIC PREPARATION CONTAINING THE SAME

**GERMAN-TITLE:** STABILISIERTE TRANSGLUTAMINASE UND DIESE ENTHALTENDE ENZYMATISCHE ZUBEREITUNG

**FRENCH-TITLE:** TRANSGLUTAMINASE STABILISEE ET PREPARATION ENZYMATIQUE CONTENANT CETTE TRANSGLUTAMINASE

**INVENTOR:** SOEDA, Takahiko Food Rch & Labs, Ajinomoto Co. Inc - 1-1, Suzuki-cho Kawasaki-ku, Kawasaki-shi, Kanagawa-ken 210, Japan (JP); HONDO, Keiko Food Rch & Labs, Ajinomoto Co., Inc. - 1-1, Suzuki-cho Kawasaki-ku, Kawasaki-shi, Kanagawa-ken 210, Japan (JP); ISHII, Chiho Food Rch & Labs, Ajinomoto Co., Inc. - 1-1, Suzuki-cho, Kawasaki-ku, Kanagawa-ken 210, Japan (JP)

**APPL-NO:** 95934272

**DESIGNATED STATES:** Belgium (BE), Germany (DE), Spain (ES), France (FR), Great Britain (GB), Italy (IT), Netherlands (NL)

**FILED-DATE:** October 11, 1995

**GRANTED-DATE:** June 23, 2004

**PRIORITY:** October 11, 1994 - 24521194, Japan (JP)

**ASSIGNEE-AT-ISSUE:** Ajinomoto Co., Inc., 00201191, No. 15-1, Kyobashi 1- chome, Chuo-ku, Tokyo 104, Japan (JP)

**LEGAL-REP:** Strehl Schubel-Hopf & Partner - 00100941, Maximilianstrasse 54, 80538 Munchen

**PUB-TYPE:** June 15, 2005 - Granted patent (B1)

**PUB-COUNTRY:** European Patent Office (EP)

**BULLETIN ISSUE NO:** June 15, 2005 - Granted patent (B1) , Issue No.: 200524

**LEGAL-STATUS:**

June 19, 1996 - Date of filing of request for examination; February 18, 2002 - Date of dispatch of the first examination report

**FILING-LANG:** Japanese (JA) (JPN)

**PUB-LANG:** English (EN) (ENG)

**PROC-LANG:** English (EN) (ENG)

**PCT-FILED-DATE:** October 11, 1995

**PCT-APPL-NO:** JP1995002076

**PCT-PUB-PAT-NO:** WO 19/96011264

**PCT-PUB-DATE:** April 18, 1996

**IPC-MAIN-CL:** 7C 12N009#96

**IPC-ADDL-CL:** C 12N009#10

**SECOND-PUBLISH:** June 23, 2004, Issue No.: 200426

**REF-CITED:**

5156956, United States (US)  
0520748, European Patent Office (EP)  
0090886, European Patent Office (EP)  
0450898, European Patent Office (EP)  
0733702, European Patent Office (EP)  
4207194, Japan (JP)

**NON-PATENT LITERATURE:** PATENT ABSTRACTS OF JAPAN vol. 016, no. 544 (C-1004), 13 November 1992 (1992-11-13) & JP 04 207194 A (AJINOMOTO KK (JP); AMANO PHARM KK (JP)), 29 July 1992 (1992-07-29)  
See also references of WO 9611264A1

**CORE TERMS:** protein, preparation, partial, hydrolysate, enzyme, aforementioned, stabilizing, transglutaminase, stabilized, sample, milk, mixture, drying, mixed, manufactured, mfd, container, minutes, peptide, wheat, stabilization, subjected, excellent, oxygen, sodium, yogurt, acid, food, dry, ice cream

**ENGLISH-ABST:**

This invention relates to stabilized transglutaminase which is obtained by drying a solution containing transglutaminase and a protein material and to a transglutaminase enzyme preparation that contains the stabilized transglutaminase as an active ingredient, wherein a partial protein hydrolysate is preferred as the protein material.

**DETDESC:**

This invention relates to a stable transglutaminase preparation having excellent shelf life (as the case may be, transglutaminase is abbreviated as TGase hereinafter).  
(Background Art)

Decreased enzyme activities are observable in many enzymes when stored for a prolonged period of time.

TGase is an enzyme whose enzyme activity is decreased in a marked degree due, mainly, to oxygen when stored for a prolonged period of time. In consequence, a process in which an organic acid, an inorganic acid, a polyphenol, a thiol compound, a sugar alcohol and the like are used as additives has been developed with the aim of improving shelf life of TGase (cf. Japanese Patent Application JP 4 207 194 A). However, improvement of shelf life by these additives is not sufficient. Especially, when TGase is stored under severe conditions such as storage throughout the summer season after its production, it is necessary to provide a proper means such as the use of an oxygen scavenger or vacuum treatment of the packaging, but these means are also still unsatisfactory.  
(Disclosure of the Invention)

In view of the above, it therefore becomes an object of the present invention to provide a TGase enzyme preparation which can be stored for a prolonged period of time at ordinary temperature without requiring oxygen scavenger, vacuum packaging and the like and a transglutaminase enzyme preparation which contains the same.

With the aim of achieving the aforementioned object, the inventors of the present invention have conducted intensive studies and found that treatment of TGase by a specified method using a specified substance can stabilize it markedly so that its enzyme activity does not decrease even after a prolonged period of storage at ordinary temperature, and have accomplished the present invention on the basis of such a finding.

Accordingly, the present invention relates to a TGase enzyme preparation obtainable by drying a solution containing TGase and a partial protein hydrolysate.

The following illustratively describes the present invention.

Origin of the TGase to be stabilized in the present invention is not particularly limited, with the proviso that it is an enzyme having TGase activity. Examples of such an enzyme include those which are originated from mammals such as guinea pig and the like (cf. Japanese Patent Publication No. JP 1 050 382 B) microorganisms such as of the genera *Streptovorticillium* and the like (for example, see Japanese Patent Application JP 1 027 471 A) and fishes such as codfish and the like (for example, see Nobuo Seki *et al.*, *Japan Journal of Scientific Fisheries*, vol.56, p.125, 1990) and those which are obtained by recombinant DNA techniques making use of biotechnology (cf. Japanese Patent Application JP 1 300 889 A, JP 5 199 883 A and JP 6 225 775 A). Of these enzymes, microbial TGase is desirable from the view point that it can be produced in a large scale and does not require calcium for the expression of its enzyme activity.

Purity of the TGase to be stabilized also has no particular limitation. That is, either crude product or high purity product obtained by purification can be stabilized.

Next, a partial protein hydrolysate to be used as a stabilizing agent is described.

The partial protein hydrolysate can be obtained in the usual way. That is, using papain, bromelain, trypsin or the like enzyme, hydrochloric acid or the like acid or sodium hydroxide or the like alkali as a partial hydrolysis agent, a protein material is partially hydrolyzed under

usually used partial hydrolysis conditions. As a matter of course, commercially available partial hydrolysates may also be used. The degree of hydrolysis of the partial hydrolysate has no particular limitation, provided that it has the TGase stabilizing function. In addition, the partial protein hydrolysate to be used in the present invention may be not only a purified product but also a product partially contaminated with impurities such as amino acids formed during the protein hydrolysis, provided that it has the TGase stabilizing function.

Drying of a solution containing TGase and a partial protein hydrolysate, namely stabilization of TGase by the aforementioned specified method using a partial protein hydrolysate, may be effected for example by either method in which (1) a partial protein hydrolysate is added during the production process of TGase or (2) a partial protein hydrolysate as the stabilizing agent is added to the produced TGase powder in a solvent, followed by dispersion and mixing and subsequent drying.

Illustratively, in the case of the method (1) in which a partial protein hydrolysate is added during the production process of TGase, a partial protein hydrolysate as the stabilizing agent is added to an enzyme solution sterilized after completion of culturing for the production of TGase in a production process disclosed for example in Japanese Patent Application JP 1 027 471 A (corresponding U.S. Patent 5,156,956), immediately or during its concentration step, subsequently drying the mixture. Of the two adding modes, the latter case in which a partial protein hydrolysate is added to the enzyme solution after its concentration to a certain level is preferred, because the partial protein hydrolysate can be dissolved easily. Particularly, wheat protein and the like can be dissolved and uniformed more easily when a TGase solution becomes viscous by its concentration to a certain level.

When the method (2) is employed, the produced TGase powder and a partial protein hydrolysate are dispersed in a solvent and mixed. This case is a method in which a partial protein hydrolysate as the stabilizing agent and TGase are uniformly mixed by dispersing or dissolving them in a solvent such as water, a buffer solution, ethanol or the like and then dried.

In both cases of the methods (1) and (2), it is not necessary to positively set a retention time after the addition of the partial protein hydrolysate and before the drying, but it is desirable to set a retention time within such a period that the enzyme activity does not decrease.

In this connection, the drying can be effected by a usually used means such as drying under reduced pressure, freeze drying or the like which does not cause inactivation of TGase.

Next, amounts of the TGase and stabilizing agent to be used when TGase is stabilized by the present invention are described. In the stabilizing methods (1) and (2) described above, there is no difference when these amounts are expressed by converting them to dry basis. That is, when TGase has a specific activity of approximately from 0.1 to 10 units/mg, the protein material is used in an adding ratio of from 0.01 to 200 weight parts, preferably from 0.5 to 100 weight parts, per 1 weight part of the enzyme. When the amount of the partial protein hydrolysate is less than 1 weight part, its function to protect TGase is so weak that stabilization of the enzyme cannot be achieved. On the other hand, when the amount exceeds 200 weight parts, its protection function can be obtained but the resulting stabilized TGase has such a low activity that it is not suitable for practical use in food due to low expression of the TGase function.

A point which should not be overlooked in relation to the present invention is that TGase cannot be stabilized when dry powder of TGase is simply mixed with dry powder of a protein material even if TGase and the protein material are used within the aforementioned range. Stabilization of TGase can be attained for the first time by drying a solution containing TGase and the partial protein hydrolysate. By doing this, it seems that TGase is incorporated into the matrix of a partial protein hydrolysate, particularly into the matrix of the partial protein

hydrolysate in liquid form in which crosslink bonding (to be described later) is formed by TGase, and, as the result, protected from oxygen in the air and stabilized.

The transglutaminase preparation (dry form) obtained in this manner may be used or distributed as it is, or other components (dry form) may be added to the transglutaminase preparation (dry form) obtained in this manner and used or distributed in the form of an enzyme preparation.

Though it varies depending on the use of the enzyme preparation of the present invention, the component to be added may be selected optionally from additives generally used in enzyme preparations or used in the prior art TGase enzyme preparation. As such additives, various protein materials may be used, including milk proteins such as powdered milk, casein, casein sodium, casein calcium and the like, as well as wheat protein, soybean protein and the like. In addition, polyphenol, organic acid salts such as citrate and the like may also be blended. Also can be blended are seasoning materials such as salt, pepper, sugar, sodium glutamate, sodium inosinate, sodium guanylate and the like and emulsifiers such as lecithin, monoglyceride and the like.

In this connection, the TGase content of the enzyme preparation of the present invention is generally from 0.1 to 99 weight parts, preferably from 1 to 90 weight parts, per 100 weight parts of the preparation. The content if less than 0.1 weight part would bear no function as the enzyme even with considerably increased dose of the enzyme preparation and if exceeding 99 weight parts would pose a difficulty in handling the preparation, because its amount to be added to the food of interest would become extremely small.

In comparison with those which are not stabilized, the TGase preparation of the present invention shows markedly small degree of periodical decrease in the enzyme activity and maintains at least about 80% of the initial activity even after one year of its production, so that it can be used effectively in a broad range of applications which include modification and adhesion of dairy products such as ice cream, yogurt, cheese and the like, animal or fish paste products such as ham, sausage, boiled fish paste and the like and various types of meat such as beef, pork, chicken meat and the like, as well as of baked wheat products such as bread and the like and pastas such as spaghetti and the like. Such effects are obtained by the modification of protein and peptide contained in materials of these products through the formation of intramolecular or intermolecular crosslink bonding generated by the enzyme action of TGase.

Though the TGase preparation obtained by the present invention has markedly high stability and shows extremely small degree of periodical inactivation as described in the foregoing, the degree of periodical inactivation can be further reduced by preserving or distributing it in an oxygen impermeable container having low oxygen permeability. It is desirable that the material of such a container has an oxygen permeability of 100 ml/m (2)[middot] 101325 Pa [middot]24h or less, preferably 30 ml/m (2)[middot]atm[middot]24 h or less. Examples of the container which satisfies the above criteria include containers in the form of a bag, a bottle and other shapes made of aluminum foil, aluminum deposition film, a polyvinyl alcohol resin, a nylon resin, a polyvinylidene chloride resin and the like.

In this connection, the activity unit of TGase is measured and defined as follows. That is, the enzyme reaction is carried out using benzyloxycarbonyl-L-glutaminyglycine and hydroxylamine as substrates, the thus formed hydroxamic acid is allowed to form its iron complex in the presence of trichloroacetic acid and then absorbance at 525 nm is measured to determine the amount of hydroxamic acid using a calibration curve and calculate the enzyme activity (cf. aforementioned Japanese Patent Application JP 1 027 471 A).

As it is universally known, TGase is an enzyme which catalyzes acyl rearrangement reaction of the gamma-carboxyamide group of glutamine residues in peptide chains. When the [epsi]-

amino group of lysine residues in protein acts as an acyl receptor, TGase catalyzes formation of [epsi]-( $\gamma$ -Glu)-Lys crosslink bonding in or between the protein molecules. Making use of such a function of TGase, modification of protein or peptide can be made (cf. Use Examples 1 to 5 which will be described later). Also, when water functions as an acyl receptor, this enzyme promotes a reaction in which glutamine residue is deaminated and converted into glutamic acid residue.

As has been described in the foregoing, the present invention relates to a TGase preparation obtainable by drying a solution containing TGase and a partial protein hydrolysate. Since the thus obtained TGase preparation is protected from the reduction of its enzyme activity caused by oxidation, it has a great advantage in that it can be preserved stably for a prolonged period of time at ordinary temperature in an oxygen-containing atmosphere.

As a matter of course, it is evident that the TGase preparation of the present invention can be provided with more higher stability, not only by containing them in oxygen-impermeable containers as described in the foregoing but also by further adding an oxygen scavenger thereto or by employing vacuum packaging or nitrogen- filled packaging.  
(Best Mode of Carrying out the Invention)

The following describes the present invention further in detail with reference to some examples. As a matter of course, the technical range of the present invention is not limited by these examples.

#### Inventive Example 1

Using a Hovert[R] Mixer (manufactured by Bokusui), 1 kg of microbial TGase (specific activity, 1.0 unit/mg) prepared in accordance with the method of Example 1 disclosed in the aforementioned Japanese Patent Application JP 1 027 471 A was thoroughly mixed with 1 kg of each of the partial protein hydrolysates shown in the following Table 1 or, for the sake of comparison, with 10 g of sodium citrate disclosed in the aforementioned Japanese Patent Application JP 4 207 194 A (control (2)), and the resulting mixture was added to 5 kg of water. After dissolving the components by mixing and allowing the resulting solution to stand at room temperature for about 30 minutes, TGase was treated by drying the solution under a reduced pressure. In each case, specific activity after the treatment was around 0.45 unit/mg.

The thus treated TGase preparation was packed and sealed in a container made of an aluminum laminate film (PET 12 [micro]/PE 15 [micro]/Al 9 [micro]/PE 15 [micro]/L-LDPE 70 [micro]) and preserved at 24[deg]C for 1 year. Results of the measurement of activities of the preserved TGase preparation samples are also shown in Table 1. In this case, a TGase sample which has been treated without adding a partial protein hydrolysate and the like stabilizing agents was also prepared to be used as a control (control (1)). Also, a powder mixture consisting of 1 kg of the same transglutaminase having a specific activity of 1.0 unit/mg and 1 kg of "Sun Lacto"[R] was preserved under the same conditions to be used as a control (control (3)).

As shown in Table 1, it was confirmed that each of the TGase samples which have been subjected to the stabilization treatment of the present invention using a partial hydrolysate as the stabilizing agent was stabilized with a significant effect, in comparison with the controls. Also, the partial protein hydrolysate was significantly excellent even in comparison with sodium citrate usually used as a stabilizing agent. In addition to the above, stabilizing effect of the partial protein hydrolysate was not found when used as a powder mixture.

Table contents not available. Please see the table in the original document. The original document or individual tables may be available via links from this document.

#### Inventive Example 2

A fermentation broth of TGase prepared in accordance with the method of Example 1 disclosed in the aforementioned Japanese Patent Application JP 1 027 471 A was filtered, the resulting filtrate was subjected to ethanol separation to obtain a precipitate (specific activity, 0.5 unit/mg) and then 1 weight part of each of the stabilizing agents shown in Table 2 was added to 1 solid weight part of the thus obtained precipitate and uniformly mixed using a mixer. This was then dried under a reduced pressure to obtain the TGase preparation of the present invention. In each case, specific activity was around 1.5 unit/mg.

The thus treated TGase preparation was packed and sealed in the same aluminum laminate film container of Inventive Example 1 and preserved at 24[deg]C for 1 year. Results of the measurement of activities of the preserved enzyme samples are also shown in Table 2. In this case, a TGase sample which has been treated without adding a partial protein hydrolysate as the stabilizing agent was also prepared to be used as a control.

As shown in Table 2, it was confirmed that each of the TGase samples which have been subjected to the stabilization treatment of the present invention was stabilized with a significant effect, in comparison with the control. The enzyme activity just after drying of each sample subjected to the treatment of the present invention is also shown in the same table, and the results also confirm that loss of the enzyme activity during the stabilization treatment of the enzyme can be reduced by adding the stabilizing agent.

Table contents not available. Please see the table in the original document. The original document or individual tables may be available via links from this document.

From the **transglutaminase** preparation samples of Inventive Example 2 which have been stored for one year after the preparation, the sample stabilized using a wheat protein partial hydrolysate ("Glutamine Peptide"[R] manufactured by DMV Japan) was selected and added to various foods described below to evaluate enzymatic function of **transglutaminase**. In this case, the **transglutaminase** preparation stabilized using "Glutamine Peptide"[R] is used herein simply as a typical example, and it is a matter of course that **transglutaminase** preparation samples prepared using other stabilizing agents than the "Glutamine Peptide"[R] can also show similar function. When the TGase preparation of the present invention is allowed to act upon protein or peptide, [epsilon]-( $\gamma$ -Glu)-Lys **crosslink** bonds are formed in or between protein or peptide molecules, thereby rendering possible modification of protein or peptide.

Use Example 1 (Meat sausage)

To 1,000 g of 3 mm square pork having a fat content of about 30% were added 20 g of salt and 450 g of ice water, followed by thorough mixing using a Stefan cutter. To this were further added 70 g of a soybean protein article ("Ajipron S2"[R] manufactured by Ajinomoto), 5 g of polyphosphate, 10 g of a seasoning ("I-7"[R] manufactured by Ajinomoto) and 10 g (15,000 units) of the aforementioned TGase preparation, subsequently mixing the contents using a Stefan cutter until final temperature of the mixture became 10[deg]C or below. In this case, the amount of added TGase was 2.5 units per 1 g protein. The thus obtained meat paste was packed in a casing, subjected to 15 minutes of smoking at 80[deg]C in a smoke house and 45 minutes of boiling at 75[deg]C and then cooled to obtain a frankfurter.

As controls, two products were prepared by repeating the same procedure except that the aforementioned TGase preparation was not added in one control and the aforementioned TGase preparation was not added and the amount of ice water to be added was reduced to 250 g in the other control.

When the thus obtained three types of frankfurters were evaluated by comparing them organoleptically, the control in which 450 g of ice water was added but TGase was not used

showed a poor touch to the teeth with no elasticity. Also, its shape as the final product became lean. However, when TGase was used, the product prepared using 450 g of ice water showed excellent quality, maintaining its elasticity and shape which were similar or superior to those of a product prepared using 250 g of ice water. Thus, the use of the aforementioned TGase preparation has rendered possible further water thinning (increase in water addition ratio) while keeping the product quality in the production of sausages.  
Use Example 2 (Ice cream)

A 200 g portion of milk was put into a 1 liter capacity container and warmed at 50[deg]C on a low fire, to which was subsequently added thoroughly loosened three egg yolks. To this was added 0.2 g the aforementioned TGase preparation (1 unit per 1 g protein of the mixture), subsequently keeping the resulting mixture for 10 minutes. During this period, temperature of the mixture was controlled at 40 to 50[deg]C. Next, 150 g of sugar and a table spoon-full of corn starch were added thereto and thoroughly mixed and then the mixture was heated on a low fire while thoroughly mixing until it becomes thick. Next, the bottom of the container was put into ice water and a few drops of vanilla essence was added while mixing the contents of the container. A 150 g portion of raw cream was added thereto and thoroughly mixed. Next, the thus obtained ice cream mix was transferred into a freezer and kept for 1 to 2 hours until its surface became solid, and then the entire portion was mixed. By repeating this step three times, an ice cream was obtained. In this case, for the sake of comparison, an ice cream (control product) was prepared by repeating the same process except that TGase was not added.

When the thus obtained two types of ice creams were organoleptically evaluated, the ice cream prepared in accordance with the present invention was excellent in spoon handling, smooth, strong in oily feeling and excellent in the taste and flavor. Also, its shape-keeping ability after 20 minutes of standing at room temperature was excellent in comparison with the control product, keeping its original shape almost completely though a slight stickiness was observed. These characteristic properties can be regarded as a result of the stabilization of milk protein emulsion effected by the action of TGase upon milk protein to form Glu-Lys bonding.

Use Example 3 (Yogurt)

A 1 kg portion of milk (fat-free solid content, 8.3%) was put into a 2 liter capacity beaker and mixed with the aforementioned preparation TGase in a ratio of 1 unit per 1 g milk protein, and the mixture was stirred for 1 hour at 25[deg]C. After 1 hour of standing, this was heated at 95[deg]C to inactivate transglutaminase. When the temperature reached 95 [deg]C, this was immediately cooled down to 44[deg]C and mixed with 1.5% per milk material of a commercial lactic acid bacterium starter "Joghurt V2"[R] (manufactured by WIESY) to carry out about 4.5 hours of fermentation at the same temperature (inventive product). The thus obtained yogurt trial product showed a pH value of 4.65. For the sake of comparison, a yogurt sample was prepared for trial by repeating the same process except that the aforementioned TGase preparation was not used (control product).

When each of these trial products was allowed to stand for 3 weeks in a refrigerator (about 15[deg]C), water separation ratio (weight ratio of separated milk serum to total yogurt) of the TGase- free control product reached 4.5%, but the TGase-added inventive product showed only 0.2% of the separation ratio, hardly showing water separation in appearance when observed by the naked eye, and the effect of TGase was still significant even when 40 days passed after its production.

According to an organoleptic evaluation carried out by 20 expert panelists 40 days after the production, the control product contained some raw granules and its eating touch was rough, while raw granules were not found in the inventive product which showed smooth eating touch inherent to yogurt. Similar to the aforementioned case, these characteristic properties can be regarded as a result of the stabilization of milk protein emulsification effected by the



action of TGase upon milk protein to form Glu-Lys bonding.  
Use Example 4 (Bread)

A 700 g portion of wheat flour (first grade hard flour) was mixed with 20 g of yeast, 1.3 g of yeast food and 380 g of water using a mixer, and the mixture was allowed to stand for 4 hours at 25[deg]C and at a humidity of 75% to effect primary fermentation, thereby obtaining a sponge dough. At the time of the completion of the fermentation, the dough showed a temperature of 28[deg]C and a pH value of 5.3. Using a mixer, this sponge dough was further kneaded with 300 g of other dough material (first grade semi-hard wheat flour), 20 g of salt, 30 g of sugar, 30 g of glucose, 30 g of shortening, 220 g of water and 0.2 unit per 1 g protein of the aforementioned TGase preparation. After allowing to stand for about 10 minutes at the same temperature of the kneading, this was equally divided into 5 balls, allowed to stand for 10 minutes at 28[deg]C and then packed in a baking mold. This was again subjected to 50 minutes of fermentation at 37[deg]C and at a humidity of 85%. Thereafter, the thus fermented dough was put into an oven and baked at 220[deg]C for 40 minutes to prepare white bread (inventive product). As a control, another white bread was prepared by repeating the same process except that the aforementioned TGase preparation was not used (control product).

The thus obtained two types of white bread samples were subjected to organoleptic evaluation in the following manner. That is, each of the white bread samples 4 days after the baking was cut in slices of 1.5 cm in thickness to carry out absolute organoleptic judgement by 10 panelists (5 males and 5 females) in terms of its texture, appearance, elasticity and eating touch, thereby finding that the product was excellent in all of the evaluation items in comparison with those of the TGase-free control product.

Use Example 5 (Spaghetti)

A 2,000 g portion of durum semolina flour ("Leone B"[R] manufactured by Nisshin Flour Milling) was mixed with 1 unit per 1 g flour protein of the aforementioned TGase preparation and 600 g of city water, and the mixture was kneaded for 10 minutes and immediately subjected to an extrusion noodle making to obtain spaghetti cut to about 40 cm in length (inventive product). At the same time, another spaghetti was prepared by repeating the same process except that the aforementioned TGase preparation was not used (control product).

Results of the evaluation of the thus obtained noodle samples showed that the TGase-added product was excellent in hardness and elasticity, namely visco-elasticity, and organoleptically desirable, in comparison with the TGase-free control product.  
(Industrial Applicability)

According to the present invention, a TGase preparation which shows extremely low level of activity loss even after standing at ordinary temperature can be obtained easily. This transglutaminase enzyme preparation of the present invention can be stored at ordinary temperature for a prolonged period of time.

In addition, not only the TGase enzyme preparation of the present invention can be used in the production of various foods such as sausage, ice cream, yogurt, bread, spaghetti and the like, but also proteins or peptides contained in raw materials of these foods can be modified by the use thereof and the quality of these foods can therefore be improved thereby.

#### **ENGLISH-CLAIMS:**

[Return to Top of Patent](#)

- A **transglutaminase** preparation comprising a **transglutaminase** enzyme and a partial protein hydrolysate, which is obtainable by drying a solution containing **transglutaminase** and the partial protein hydrolysate.

- The **transglutaminase** preparation according to claim 1 which is in freeze-dried form.

- A **transglutaminase** enzyme preparation which comprises the stabilized **transglutaminase** preparation of claim 1 or 2 and an additive agent.
- The **transglutaminase** enzyme preparation according to claim 3, wherein the additive agent is at least one member selected from polyphenol and organic acid salts.
- A method for modifying proteins or peptides which comprises allowing the stabilized **transglutaminase** preparation of claim 1 or 2 or the enzyme preparation of claim 3 or 4 to act upon a protein or a peptide, thereby effecting formation of [epsi]-(gamma-Glu)-Lys **crosslink** bonding in or between molecules of the protein or peptide.
- A **transglutaminase** package in which the stabilized **transglutaminase** preparation of claim 1 or 2 or the enzyme preparation of claim 3 or 4 is packed and sealed in a container made of a material having an oxygen permeability of 100 ml/m (2)[middot]101325 Pa [middot]24h or less or subjected to vacuum packaging or nitrogen-filled packaging optionally together with an oxygen scavenger.

#### **GERMAN-CLAIMS:**

- Transglutaminasezubereitung, umfassend eine Transglutaminaseenzym und ein Teilproteinhydrolysat, erhaltlich durch Trocknen einer Losung, die Transglutaminase und ein Teilproteinhydrolysat enthalt.
- 'Transglutaminasezubereitung nach Anspruch 1, die in gefriergetrockneter Form vorliegt.
- Transglutaminaseenzymzubereitung, welche die stabilisierte Transglutaminasezubereitung nach Anspruch 1 oder 2 und ein Zusatzmittel umfast.
- Transglutaminaseenzymzubereitung nach Anspruch 3, worin das Zusatzmittel mindestens ein Mitglied ist, ausgewählt aus Polyphenol und Salzen organischer Sauren.
- Verfahren zur Modifizierung von Proteinen oder Peptiden, umfassend das Ermoglichen des Einwirkens der stabilisierten Transglutaminasezubereitung nach Anspruch 1 oder 2 oder deren Enzymzubereitung der Anspruche 3 oder 4 auf ein Protein oder ein Peptid, wodurch die Bildung einer [epsi]-(gamma -Glu)-Lys-Vernetzungsbindung in oder zwischen Molekullen des Protein oder Peptids bewirkt wird.
- Transglutaminasepackung, in welcher die stabilisierte Transglutaminasezubereitung nach Anspruch 1 oder 2 oder die Enzymzubereitung von Anspruch 3 oder 4 in einem Behalter aus einem Material mit einer Sauerstoffdurchlassigkeit von 100 ml/m ( 2)[middot]101325 Pa [middot]24 h oder weniger verpackt oder versiegelt ist, oder einer Vakuumverpackung unterzogen wurde, oder einer Stickstoff-gefullten Verpackung, normalerweise zusammen mit einem Sauerstofffanger.

#### **FRENCH-CLAIMS:**

- Preparation de transglutaminase comprenant l'enzyme transglutaminase et un hydrolysat de proteine partielle qui peut etre obtenue par sechage d'une solution contenant la transglutaminase et l'hydrolysat de proteine partielle.
- Preparation de transglutaminase selon la revendication 1, qui se presente sous forme lyophilisee.
- Preparation enzymatique de transglutaminase qui comprend la preparation de transglutaminase stabilisee selon la revendication 1 ou la revendication 2 et un additif.
- Preparation enzymatique de transglutaminase selon la revendication 3, dans laquelle l'additif est au moins un membre choisi parmi le polyphenol et des sels d'acides organiques.
- Procede de modification de proteines ou de peptides qui comprend le fait de laisser la preparation de transglutaminase stabilisee selon la revendication 1 ou la revendication 2 ou la preparation enzymatique selon la revendication 3 ou la revendication 4 pour agir sur une proteine ou un peptide, effectuant ainsi la formation de liaison de reticulation [epsi]-(gamma-Glu) Lys dans ou entre les molecules de la proteine ou du peptide.
- Emballage pour transglutaminase dans lequel la preparation de transglutaminase stabilisee selon la revendication 1 ou la revendication 2 ou la preparation enzymatique selon la revendication 3 ou la revendication 4 est emballee et scellee dans un recipient fabrique dans un materiau ayant une permeabilite a l'oxygene de 100 ml/m (2) . 101325 Pa . 24 h ou

moins ou soumis a un emballage sous vide ou un emballage a atmosphere d'azote  
eventuellement en association a un eliminateur d'oxygene.

Source: [Legal](#) > / . . . / > **U.S. Patents, European Patents, Patent Abstracts of Japan, PCT Patents, and U.K.** 

Terms: **transglutaminase /p (cross-link! or crosslink!) and date<8/1/1998** ([Edit Search](#) | [Suggest Terms for My Search](#))

View: Full

Date/Time: Friday, May 12, 2006 - 8:15 PM EDT



[About LexisNexis](#) | [Terms & Conditions](#)

[Copyright](#) © 2006 LexisNexis, a division of Reed Elsevier Inc. All rights reserved.